

Finite orbits for rational functions

Jung Kyu CANCI

Abstract

Let K be a number field and $\phi \in K(z)$ a rational function. Let S be the set of all archimedean places of K and all non-archimedean places associated to the prime ideals of bad reduction for ϕ . We prove an upper bound for the length of finite orbits for ϕ in $\mathbb{P}_1(K)$ depending only on the cardinality of S .

Introduction

Let K be a number field and R its ring of integers. With every rational function $\phi \in K(z)$ we associate in the canonical way a rational map $\Phi: \mathbb{P}_1 \rightarrow \mathbb{P}_1$ defined over K . For every point $P \in \mathbb{P}_1(K)$ we call its *forward orbit under Φ* (or simply *orbit*) the set $O_\Phi(P) = \{\Phi^n(P) \mid n \in \mathbb{Z}_{\geq 0}\}$, where Φ^n is the n -th iterate of Φ and $\Phi^0(P) = P$. If $O_\Phi(P)$ is a finite set one says that P is a *pre-periodic* point for Φ . This definition is due to the following fact: if $O_\Phi(P)$ is finite then there exist two integers $n \in \mathbb{Z}_{\geq 0}$ and $m \in \mathbb{Z}_{> 0}$ such that $\Phi^n(P) = \Phi^{n+m}(P)$. In this case one says that $\Phi^n(P)$ is a *periodic* point for Φ . If m is the smallest positive integer with the above property, then one says that m is the *period* of P . If P is a periodic point then its orbit is called a *cycle*.

It is not difficult to prove that every polynomial in $\mathbb{Z}[x]$ has cycles in $\mathbb{Z}$ of length at most 2 and every finite orbit has cardinality at most 6. For a fixed finite set S of valuations of K , containing all the archimedean ones, Narkiewicz in [11] has shown that if Φ is a monic polynomial with coefficients in the ring of S -integers R_S (see the definition at the beginning of the next section), then the length of its cycles in K is bounded by a function $B(R_S) = C^{|S|(|S|+2)}$, for an absolute constant C . Note that the bound depends only on the cardinality of S . The value of $B(R_S)$ has been diminished by Pezda in [13]. Indeed, the main result of Pezda [13, Theorem 1], which concerns polynomial maps in local rings, combined with the estimate given in [1, Theorem 4.7] on the height of the $|S|$ -th rational prime,

2000 Mathematics Subject Classification: 11G99, 14E05

Key words: rational maps, finite orbits, S -unit equations, reduction modulo $\mathfrak{p}$

gives rise to the following inequality

$$B(R_S) \leq [12|S| \log(5|S|)]^{2[K:\mathbb{Q}]+1}. \quad (1)$$

Later Narkiewicz and Pezda in [12] extended [13, Theorem 1] to finite orbits so including pre-periodic points. By considering the limit in (1) and the Evertse's bound proved in [6] for the number of S -unit non-degenerate solutions to linear equations in three variables, the result of Narkiewicz and Pezda [12, Theorem 1] states that the length of a finite orbit in K for a monic polynomial with coefficients in R_S is at most

$$\frac{1}{3} [12|S| \log(5|S|)]^{2[K:\mathbb{Q}]+1} (31 + 2^{1031|S|}) - 1.$$

R. Benedetto has recently obtained a different bound, again for polynomial maps, but his bound also depends on the degree of the map. He proved in [2] that if $\phi \in K[z]$ is a polynomial of degree $d \geq 2$ which has bad reduction at s primes of K , then the number of pre-periodic points of ϕ is at most $O(s \log s)$. The big- O constant is essentially $(d^2 - 2d + 2)/\log d$ for large s . Benedetto's proof relies on a detailed analysis of p -adic Julia sets.

In the present paper we will generalize to finite orbits for rational maps the result of Narkiewicz and Pezda [12] obtained for polynomial maps. We will study the same semigroup of rational maps studied in [4], namely: we fix an arbitrary finite set S of places of K containing all archimedean ones and consider the rational maps with good reduction outside S . We recall the definition of good reduction for a rational map at a non zero prime ideal $\mathfrak{p}$ (for the details see [10] or [4]): a rational map $\Phi: \mathbb{P}_1 \rightarrow \mathbb{P}_1$, defined over K , has good reduction at a prime ideal $\mathfrak{p}$ if there exists a rational map $\tilde{\Phi}: \mathbb{P}_1 \rightarrow \mathbb{P}_1$, defined over $K(\mathfrak{p})$, such that $\deg \Phi = \deg \tilde{\Phi}$ and the following diagram

$$\begin{array}{ccc} \mathbb{P}_{1,K} & \xrightarrow{\Phi} & \mathbb{P}_{1,K} \\ \downarrow \sim & & \downarrow \sim \\ \mathbb{P}_{1,K(\mathfrak{p})} & \xrightarrow{\tilde{\Phi}} & \mathbb{P}_{1,K(\mathfrak{p})} \end{array}$$

is commutative, where $\sim$ is the reduction modulo $\mathfrak{p}$. In other words, an endomorphism Φ of $\mathbb{P}_1$ defined over K has good reduction at $\mathfrak{p}$ if Φ can be written as $\Phi([X : Y]) = [F(X, Y), G(X, Y)]$, where F and G are homogeneous polynomials of the same degree, with coefficients in the local ring $R_{\mathfrak{p}}$ of R at $\mathfrak{p}$, and such that the resultant $\text{Res}(F, G)$ of polynomials F and G is a $\mathfrak{p}$ -unit in $R_{\mathfrak{p}}$. Note that, from this definition, a rational map on $\mathbb{P}_1(K)$ associated to a polynomial in $K[z]$ has good

reduction outside S if and only if its coefficients are S -integers and its leading coefficient is an S -unit.

In this paper we prove:

Theorem 1. *Let K be a number field. Let S be a finite set of cardinality s of places of K containing all the archimedean ones. There exists a number $c(s)$, depending only on s , such that the length of every finite orbit in $\mathbb{P}_1(K)$, for rational maps with good reduction outside S , is bounded by $c(s)$. We can choose $c(s)$ equal to*

$$\left[e^{10^{12}} (s+1)^8 (\log(5(s+1)))^8 \right]^s. \quad (2)$$

The proof of Theorem 1 uses two non-elementary facts: the first is [9, Corollary B] where Morton and Silverman proved that if Φ is a rational map of degree ≥ 2 which has bad reduction only at t prime ideals of K and $P \in \mathbb{P}_1(K)$ is a periodic point with minimal period n , then the inequality

$$n \leq [12(t+2) \log(5(t+2))]^{4[K:\mathbb{Q}]} \quad (3)$$

holds. The second one is the theorem proved by Evertse, Schlickewei and Schmidt in [7] on the number of non-degenerate solutions $(u_1, \dots, u_n) \in \Gamma$ to equation $a_1 u_1 + \dots + a_n u_n = 1$ where Γ is a given subgroup of $(\mathbb{C}^*)^n$ of finite rank and the a_i 's are given non-zero elements of K . For $n = 2$ and $a_1 = a_2 = 1$ we use the upper bound proved by Beukers and Schlickewei in [3]. The main point to obtain the estimate of Theorem 1 is the fact that the upper bounds in the theorems in [7] and [3] only depend on the rank of Γ . From Theorem 1 we easily deduce the following result concerning finite orbits for rational maps contained in a given finitely generated semigroup of endomorphisms of $\mathbb{P}_1$:

Corollary 1. *Let $\mathcal{F}$ be a finitely generated semigroup of endomorphisms of $\mathbb{P}_1$ defined over a number field K . There exists a uniform upper bound C which bounds the length of every finite orbit in $\mathbb{P}_1(K)$ for any rational map in $\mathcal{F}$. Furthermore it is possible to give an explicit bound for C in terms of a set of generators of $\mathcal{F}$.*

Acknowledgements. The present work was written during the preparation of my Ph.D. thesis supervised by professor P. Corvaja. I would like to thank him for his useful suggestions. Also, I would like to thank prof. J.H. Silverman and prof. U. Zannier for their corrections and advice on my thesis. I am grateful to the referee for useful suggestions which helped to improve Theorem 1 and, in general, the presentation of this paper.

1 Proofs

In all the present paper we will use the following notation:

K a number field;

R the ring of integers of K ;

$\mathfrak{p}$ a non zero prime ideal of R ;

$v_{\mathfrak{p}}$ the $\mathfrak{p}$ -adic valuation on R corresponding to the prime ideal $\mathfrak{p}$ (we always assume $v_{\mathfrak{p}}$ to be normalized so that $v_{\mathfrak{p}}(K^*) = \mathbb{Z}$);

S a fixed finite set of places of K of cardinality s including all archimedean places;

$R_S := \{x \in K \mid v_{\mathfrak{p}}(x) \geq 0 \text{ for every prime ideal } \mathfrak{p} \notin S\}$ the ring of S -integers;

$R_S^* := \{x \in K^* \mid v_{\mathfrak{p}}(x) = 0 \text{ for every prime ideal } \mathfrak{p} \notin S\}$ the group of S -units.

Let $P_1 = [x_1 : y_1]$ and $P_2 = [x_2 : y_2]$ be points in $\mathbb{P}_1(K)$. Using the notation of [10] we will denote by

$$\delta_{\mathfrak{p}}(P_1, P_2) = v_{\mathfrak{p}}(x_1 y_2 - x_2 y_1) - \min\{v_{\mathfrak{p}}(x_1), v_{\mathfrak{p}}(y_1)\} - \min\{v_{\mathfrak{p}}(x_2), v_{\mathfrak{p}}(y_2)\} \quad (4)$$

the $\mathfrak{p}$ -adic logarithmic distance between the points P_1, P_2 ; note that $\delta_{\mathfrak{p}}(P_1, P_2)$ is independent of the choice of the homogeneous coordinates, i.e. it is well defined. We will use the two following propositions contained in [10]:

Proposition 1. [10, Proposition 5.1]

$$\delta_{\mathfrak{p}}(P_1, P_3) \geq \min\{\delta_{\mathfrak{p}}(P_1, P_2), \delta_{\mathfrak{p}}(P_2, P_3)\}$$

for all $P_1, P_2, P_3 \in \mathbb{P}_1(K)$. □

Proposition 2. [10, Proposition 5.2] *Let $\Phi: \mathbb{P}_1(K) \rightarrow \mathbb{P}_1(K)$ be a rational map defined over K . Then*

$$\delta_{\mathfrak{p}}(\Phi(P), \Phi(Q)) \geq \delta_{\mathfrak{p}}(P, Q)$$

for all $P, Q \in \mathbb{P}_1(K)$ and all prime ideals $\mathfrak{p}$ of good reduction for Φ . □

With $(Q_{-m}, \dots, Q_0, \dots, Q_{n-1})$ we always represent a finite orbit for a rational map Ψ in which the 0-th term Q_0 is a n -th periodic point for Ψ . Moreover, for all indexes $i \geq -m$, $Q_{i+1} = \Psi(Q_i)$ holds, bearing in mind that $Q_n = Q_0$. We will use the following remark which is a direct consequence of the previous two propositions.

Remark 1. Let $(Q_{-m}, \dots, Q_0, \dots, Q_{n-1})$ be a finite orbit in $\mathbb{P}_1(K)$ for a rational map Ψ with good reduction outside S ; then for all integers a, b with $-m \leq a \leq n-1, b \geq 0, k \geq 0$ and for every prime ideal $\mathfrak{p} \notin S$

$$\delta_{\mathfrak{p}}(Q_a, Q_{a+kb}) \geq \min\{\delta_{\mathfrak{p}}(Q_a, Q_{a+b}), \delta_{\mathfrak{p}}(Q_{a+b}, Q_{a+2b}), \dots, \delta_{\mathfrak{p}}(Q_{a+(k-1)b}, Q_{a+kb})\} = \delta_{\mathfrak{p}}(Q_a, Q_{a+b}).$$

Proof. It is a direct application of the triangle inequality (Proposition 1) and Proposition 2. In fact the b -th iterate of Ψ has good reduction at every prime ideal $\mathfrak{p} \notin S$, therefore

$$\delta_{\mathfrak{p}}(Q_{a+lb}, Q_{a+(l+1)b}) = \delta_{\mathfrak{p}}(\Psi^b(Q_{a+(l-1)b}), \Psi^b(Q_{a+lb})) \geq \delta_{\mathfrak{p}}(Q_{a+(l-1)b}, Q_{a+lb})$$

for all indexes $0 < l \leq k$. □

In the first version of this paper, in Theorem 1, we proved an upper bound of the form $c(s, h)$ also depending on the class number h of the ring R_S . Indeed we worked with a set $\mathbb{S}$ of places of K containing S such that the ring $R_{\mathbb{S}}$ was a principal ideal domain. From a simple inductive argument it results that it is possible to choose $\mathbb{S}$ such that $|\mathbb{S}| \leq s + h - 1$. From some suitable applications of Proposition 1 and Proposition 2, we obtained some equations in two and three $\mathbb{S}$ -units and by using the upper bounds proved by Evertse in [5] and [6] we deduced a bound in Theorem 1. Following the useful suggestions made by the anonymous referee we shall use, instead of the classical S -unit equation theorem, the refined result of Evertse, Schlickewei and Schmidt [7] (and of Beukers and Schlickewei [3] for $n = 2$) leading to an upper bound in Theorem 1 depending only on the cardinality of S , even if R_S is not a principal ideal domain. Now we state the last two quoted theorems and then we present the referee's suggestion to use these results.

Let L be a number field. Let $(L^*)^n$ be the n -fold direct product of L^* , with coordinatewise multiplication $(x_1, \dots, x_n)(y_1, \dots, y_n) = (x_1y_1, \dots, x_ny_n)$ and exponentiation $(x_1, \dots, x_n)^t = (x_1^t, \dots, x_n^t)$. We say that a subgroup Γ of $(L^*)^n$ has rank r if Γ has a free subgroup Γ_0 of rank r such that for every $\mathbf{x} \in \Gamma$ there is $m \in \mathbb{Z}_{>0}$ with $\mathbf{x}^m \in \Gamma_0$.

Theorem A [3] *Let L be a number field and let Γ be a subgroup of $(L^*)^2$ of rank r . Then the equation*

$$x + y = 1 \quad \text{in } (x, y) \in \Gamma$$

has at most $2^{8(r+1)}$ solutions. □

Theorem B [7] *Let L be a number field, let $n \geq 3$ and let $a_1, \dots, a_n$ be non zero elements of L . Further, let Γ be a subgroup of $(L^*)^n$ of rank r . Then the equation*

$$a_1 x_1 + \dots + a_n x_n = 1 \quad \text{in } (x_1, \dots, x_n) \in \Gamma$$

has at most $e^{(6n)^{3n}(r+1)}$ solutions such that $\sum_{i \in I} a_i x_i \neq 0$ for each non empty subset $I \subset \{1, \dots, n\}$. $\square$

Let $\mathbf{a}_1, \dots, \mathbf{a}_h$ be a full system of representatives for the ideal classes of R_S . For each $i \in \{1, \dots, h\}$ there is an S -integer $\alpha_i \in R_S$ such that

$$\mathbf{a}_i^h = \alpha_i R_S. \quad (5)$$

Let L be the extension of K given by

$$L = K(\zeta, \sqrt[h]{\alpha_1}, \dots, \sqrt[h]{\alpha_h}) \quad (6)$$

where ζ is a primitive h -th root of unity. Of course if $h = 1$ then $L = K$. Let us define the following subgroups of L^*

$$\sqrt{K^*} := \{a \in L^* \mid \exists m \in \mathbb{Z}_{>0} \text{ with } a^m \in K^*\}$$

and

$$\sqrt{R_S^*} := \{a \in L^* \mid \exists m \in \mathbb{Z}_{>0} \text{ with } a^m \in R_S^*\}.$$

Let $\mathbf{S}$ denote the set of places of L lying above the places in S and denote by R_S and R_S^* the ring of $\mathbf{S}$ -integers and the group of $\mathbf{S}$ -units, respectively, in L . By definition it is clear that $R_S^* \cap \sqrt{K^*} = \sqrt{R_S^*}$ and so it follows that $\sqrt{R_S^*}$ is a subgroup of L^* of rank $s - 1$. With the just stated notation we prove the following:

Proposition 3. *Let L and $\mathbf{S}$ be as above. Let Φ be a rational map from $\mathbb{P}_1$ to $\mathbb{P}_1$ defined over L , having good reduction at all prime ideals outside $\mathbf{S}$. Let*

$$\{P_{-m}, \dots, P_{-1}, P_0\} \quad (7)$$

be a set of $m + 1$ distinct points of $\mathbb{P}_1(L)$ such that $\Phi(P_i) = P_{i+1}$ for all $i \in \{-m, \dots, -1\}$ and $\Phi(P_0) = P_0$. Further, suppose that $P_i = [x_i : y_i]$ for all indexes $i \in \{-m, \dots, 0\}$, where $x_i, y_i \in L$ such that

- (1) $x_0 = 0, y_0 = 1$;
- (2) $x_i R_S + y_i R_S = R_S$ for all indexes $i \in \{-m, \dots, 0\}$;
- (3) $x_i y_j - x_j y_i \in \sqrt{K^*}$ for any distinct indexes $i, j \in \{-m, \dots, 0\}$.

Then $m < e^{10^{12}s} - 2$.

The proof of this proposition will be a direct consequence of the following three lemmas.

Lemma 1. *With the same hypothesis of Proposition 3, let $P_{l-k}, \dots, P_{l-1}, P_l$ be distinct points of the orbit (7) such that for every prime ideal $\mathfrak{p} \notin \mathbf{S}$*

$$\delta_{\mathfrak{p}}(P_{l-i}, P_0) = \delta_{\mathfrak{p}}(P_l, P_0) \text{ for every index } 0 \leq i \leq k. \quad (8)$$

Then $k < 2^{16^s}$.

Proof. For every prime ideal $\mathfrak{p} \notin \mathbf{S}$ and for any two indexes $k \geq i > j \geq 0$ from Proposition 1 and condition (8) it follows that

$$\delta_{\mathfrak{p}}(P_{l-i}, P_{l-j}) \geq \min\{\delta_{\mathfrak{p}}(P_{l-i}, P_0), \delta_{\mathfrak{p}}(P_{l-j}, P_0)\} = \delta_{\mathfrak{p}}(P_l, P_0). \quad (9)$$

Moreover, since $P_n = P_0$ for all $n \geq 0$, by applying Remark 1 to the orbit $(P_{-m}, \dots, P_{-1}, P_0)$ with $a = l - i$, $b = i - j$ and $k = (m + 1)$, where m is the maximum integer such that $l - i + m(i - j) < 0$, it follows that

$$\begin{aligned} \delta_{\mathfrak{p}}(P_{l-i}, P_0) &\geq \min\{\delta_{\mathfrak{p}}(P_{l-i}, P_{l-j}), \delta_{\mathfrak{p}}(P_{l-j}, P_{l+i-2j}), \dots, \delta_{\mathfrak{p}}(P_{l-i+m(i-j)}, P_0)\} \\ &= \delta_{\mathfrak{p}}(P_{l-i}, P_{l-j}). \end{aligned}$$

By the last inequality, (8) and (9) we have that

$$\delta_{\mathfrak{p}}(P_{l-i}, P_{l-j}) = \delta_{\mathfrak{p}}(P_l, P_0). \quad (10)$$

Note that by condition (2)-Proposition 3

$$\delta_{\mathfrak{p}}(P_i, P_j) = v_{\mathfrak{p}}(x_i y_j - x_j y_i) \quad (11)$$

for all indexes $i, j \in \{-m, \dots, 0\}$ and every prime ideal $\mathfrak{p} \notin \mathbf{S}$. Since $P_0 = [0 : 1]$, from (3)-Proposition 3 it follows that $x_{l-i} \in \sqrt{K^*}$ and so, by (11), condition (8) is equivalent to $x_{l-i} R_{\mathbf{S}} = x_l R_{\mathbf{S}}$, for every index $0 \leq i \leq k$. Hence

$$u_{l-i} := \frac{x_{l-i}}{x_l} \in R_{\mathbf{S}}^* \cap \sqrt{K^*} = \sqrt{R_{\mathbf{S}}^*} \quad (12)$$

and $P_{l-i} = [x_l : y_{l-i}/u_{l-i}]$. Furthermore, again from (3)-Proposition 3 combined with (10) and (12) we deduce that

$$u_{l-i, l-j} := \frac{x_{l-i} y_{l-j} - x_{l-j} y_{l-i}}{x_l u_{l-i} u_{l-j}} = \frac{y_{l-j}}{u_{l-j}} - \frac{y_{l-i}}{u_{l-i}} \in R_{\mathbf{S}}^* \cap \sqrt{K^*} = \sqrt{R_{\mathbf{S}}^*} \quad (13)$$

for all distinct indexes $i, j \in \{0, \dots, k\}$. In particular, either $k \in \{0, 1\}$ or we have a system of three equations

$$\begin{cases} y_l - y_{l-1}/u_{l-1} = u_{l-1,l} \\ y_l - y_{l-i}/u_{l-i} = u_{l-i,l} \\ y_{l-1}/u_{l-1} - y_{l-i}/u_{l-i} = u_{l-i,l-1} \end{cases}. \quad (14)$$

The first one is obtained from (13) substituting $j = 0$ and $i = 1$ and the two other ones with $j = 0, j = 1$ and i an arbitrary index $k \geq i \geq 2$ (recall that $u_l = 1$).

We deduce from (14) the following linear relation:

$$u_{l-1,l} + u_{l-i,l-1} = u_{l-i,l},$$

so $(u_{l-1,l}/u_{l-i,l}, u_{l-i,l-1}/u_{l-i,l}) \in \sqrt{R_S^*} \times \sqrt{R_S^*}$ is a solution of the equation $u + v = 1$. Note that the group $\sqrt{R_S^*} \times \sqrt{R_S^*}$ has rank $2(s-1)$ therefore, by Theorem A (Beukers and Schlickewei [3]) with $\Gamma = \sqrt{R_S^*} \times \sqrt{R_S^*}$, there are at most $2^{8(2s-2+1)} = 2^{16s-8}$ possibilities for $(u_{l-1,l}/u_{l-i,l}, u_{l-i,l-1}/u_{l-i,l})$. Now from (14) it follows that

$$\frac{y_{l-i}}{u_{l-i}} = y_l - \frac{u_{l-i,l}}{u_{l-1,l}} u_{l-1,l}.$$

Thus the set of points $\{P_{l-i} = [x_l : y_{l-i}/u_{l-i}] \mid k \geq i \geq 2\}$ has cardinality bounded by 2^{16s-8} so $k \leq 2^{16s-8} + 1 < 2^{16s}$. $\square$

The next step is to prove an upper bound, which depends only on s , for the number of points P_{-i} of (7) such that $x_{-i}R_S \neq x_{-i+1}R_S$. We need two lemmas.

We say that a S -integer T is representable in *two essentially different ways* as sum of two elements of $\sqrt{R_S^*}$ if there exist

$$u_1, u_2, v_1, v_2 \in \sqrt{R_S^*} \text{ such that } \{u_1, u_2\} \neq \{v_1, v_2\} \text{ and } T = u_1 + u_2 = v_1 + v_2. \quad (15)$$

Lemma 2. *The cardinality of the set of non zero principal ideals of R_S*

$$\{T \cdot R_S \mid T \text{ satisfies (15)}\}$$

is bounded by $e^{(18)^9(3s-2)}$.

Proof. Let $T \in R_S/\{0\}$ be written as $T = u_1 + u_2 = v_1 + v_2$ which satisfies the condition in (15). Therefore the left term of equation

$$\frac{u_1}{v_1} + \frac{u_2}{v_1} - \frac{v_2}{v_1} = 1$$

has no vanishing subsums. Now, applying Theorem B (Evertse, Schlickewei and Schmidt [7]) with $n = 3$ and $\Gamma = \sqrt{R_S^*} \times \sqrt{R_S^*} \times \sqrt{R_S^*}$ we obtain that the principal ideal

$$T \cdot R_S = v_1 \left(1 + \frac{v_2}{v_1}\right) \cdot R_S$$

has at most $e^{(18)^9(3s-2)}$ possibilities. $\square$

Remark 2. By previous lemma, we can choose a set $\mathfrak{T}$ of $\mathbf{S}$ -integers, with cardinality at most $e^{(18)^9(3s-2)}$, such that every non zero $\mathbf{S}$ -integer with the property (15) is representable as uT , where $u \in R_S^*$ and $T \in \mathfrak{T}$.

Lemma 3. *With the same hypothesis of Proposition 3, if there exist five distinct points $P_{n_5} = [x_{n_5} : y_{n_5}]$, $P_{n_4} = [x_{n_4} : y_{n_4}]$, $P_{n_3} = [x_{n_3} : y_{n_3}]$, $P_{n_2} = [x_{n_2} : y_{n_2}]$, $P_{n_1} = [x_{n_1} : y_{n_1}]$ of the orbit (7), with $n_5 < n_4 < n_3 < n_2 < n_1 < 0$, then x_{n_1}/x_{n_2} is a non zero $\mathbf{S}$ -integer that is representable, in two essentially different ways, as sum of two elements of $\sqrt{R_S^*}$.*

Proof. Since $\Phi(P_0) = P_0 = [0 : 1]$, from Proposition 2, considering $\Phi^{n_i - n_j}$, $P = P_{n_j}$ and $Q = P_0$, it follows that $x_{n_j}|x_{n_i}$ in R_S for all couple of integers $j \geq i$. Therefore there exist four non zero $\mathbf{S}$ -integers T_1, T_2, T_3, T_4 such that

$$x_{n_i} = T_i x_{n_{i+1}} \text{ for all } i \in \{1, 2, 3, 4\}$$

and so for every couple of distinct indexes $1 \leq i < j \leq 5$

$$x_{n_i} = T_i \cdot \dots \cdot T_{j-1} x_{n_j}. \quad (16)$$

By Remark 1 we have that

$$\delta_p(P_{n_j}, P_0) \geq \min\{\delta_p(P_{n_j}, P_{n_i}), \delta_p(P_{n_i}, P_{2n_i - n_j}), \dots, \delta_p(P_{m \cdot n_i - (m-1)n_j}, P_0)\} = \delta_p(P_{n_i}, P_{n_j})$$

for a suitable integer m , so it follows that $(x_{n_j}y_{n_i} - x_{n_i}y_{n_j})|x_{n_j}$ and by identity (16)

$$y_{n_i} - T_i \cdot \dots \cdot T_{j-1} y_{n_j} = \frac{x_{n_j}y_{n_i} - x_{n_i}y_{n_j}}{x_{n_j}} \in R_S^* \cap \sqrt{K^*} = \sqrt{R_S^*}. \quad (17)$$

(Recall that, by conditions (1) and (3) in the hypothesis of Proposition 3, the $\mathbf{S}$ -integers $x_{n_j}y_{n_i} - x_{n_i}y_{n_j}$ and x_{n_j} belong to $\sqrt{K^*}$.) From (17) we obtain:

$$y_{n_1} - T_1 y_{n_2} = v_1, \quad (18)$$

$$y_{n_2} - T_2 y_{n_3} = v_2, \quad (19)$$

$$y_{n_1} - T_1 T_2 y_{n_3} = v_3, \quad (20)$$

$$y_{n_3} - T_3 y_{n_4} = v_4, \quad (21)$$

$$y_{n_2} - T_2 T_3 y_{n_4} = v_5, \quad (22)$$

$$y_{n_1} - T_1 T_2 T_3 y_{n_4} = v_6, \quad (23)$$

$$y_{n_2} - T_2 T_3 T_4 y_{n_5} = v_7, \quad (24)$$

$$y_{n_1} - T_1 T_2 T_3 T_4 y_{n_5} = v_8, \quad (25)$$

$$y_{n_3} - T_3 T_4 y_{n_5} = v_9, \quad (26)$$

$$y_{n_4} - T_4 y_{n_5} = v_{10}, \quad (27)$$

where $v_i \in \sqrt{R_S^*}$ for all indexes $1 \leq i \leq 10$.

From (18), (20) and (19) we obtain

$$T_1 = \frac{v_3}{v_2} - \frac{v_1}{v_2}. \quad (28)$$

From (23), (18) and (22) we obtain

$$T_1 = \frac{v_6}{v_5} - \frac{v_1}{v_5}. \quad (29)$$

From (25), (18) and (24) we obtain

$$T_1 = \frac{v_8}{v_7} - \frac{v_1}{v_7}. \quad (30)$$

Now we finish to proving that among (28), (29), (30) there exist at least two distinct representations of T_1 as sum of two elements of $\sqrt{R_S^*}$. From (24), (19) and (26) we obtain that $T_2 = \frac{v_7-v_2}{v_9}$; therefore $v_7 \neq v_2$ and so

$$\left\{ \frac{v_3}{v_2}, -\frac{v_1}{v_2} \right\} = \left\{ \frac{v_8}{v_7}, -\frac{v_1}{v_7} \right\} \Rightarrow -\frac{v_1}{v_2} = \frac{v_8}{v_7}. \quad (31)$$

From (24), (22) and (27) we obtain that $T_2 T_3 = \frac{v_7-v_5}{v_{10}}$; therefore $v_7 \neq v_5$ and so

$$\left\{ \frac{v_6}{v_5}, -\frac{v_1}{v_5} \right\} = \left\{ \frac{v_8}{v_7}, -\frac{v_1}{v_7} \right\} \Rightarrow -\frac{v_1}{v_5} = \frac{v_8}{v_7}. \quad (32)$$

From (31) and (32) it follows that

$$\left\{ \frac{v_3}{v_2}, -\frac{v_1}{v_2} \right\} = \left\{ \frac{v_6}{v_5}, -\frac{v_1}{v_5} \right\} = \left\{ \frac{v_8}{v_7}, -\frac{v_1}{v_7} \right\} \Rightarrow -\frac{v_1}{v_2} = -\frac{v_1}{v_5}.$$

But this is not possible since, from (22), (19) and (21), $T_2 = \frac{v_5-v_2}{v_4} \neq 0$ holds.

□

Proof of Proposition 3. The set $\{P_{i_r}, \dots, P_{i_1}, P_{-1}\}$ of all points P_{-j} of the orbit (7) such that $x_{-j}R_S \neq x_{-j+1}R_S$ has cardinality equal to $r+1 \leq 4 + e^{(18)^9(3s-2)}$. Indeed, if such five points do not exist we have finish; otherwise for every index $i_{r-2} < i_t \leq i_1$ we apply the previous lemma with $n_1 = -1, n_2 = i_t, n_3 = i_{r-2}, n_4 = i_{r-1}, n_5 = i_r$ obtaining that $x_{-1}x_{i_t}^{-1} = uT$ where $T \in \mathfrak{T}$ (the set chosen in Remark 2) and u is a suitable $\mathbf{S}$ -unit. Therefore

$$P_{i_t} = [x_{-1}/T : uy_{i_t}].$$

In this way we have proved that r is bounded by $3 + |\mathfrak{T}|$. Now, by Lemma 1, it is clear that it is possible to choose as upper bound for m the number

$$(4 + e^{(18)^9(3s-2)})(2^{16s} + 1) < e^{(10)^{12s}} - 2.$$

□

Proof of Theorem 1. The bound (2) holds for finite orbit length for all rational maps of degree 1, i.e. automorphisms of $\mathbb{P}_1(K)$. Indeed every pre-periodic point for a bijection is a periodic point. Thus we have to study only the cycle lengths. If a point of $\mathbb{P}_1(K)$ is a periodic point for an automorphism $\Psi \in \text{PGL}_2(K)$, with period $n \geq 3$, then Ψ^n is the identity map of $\mathbb{P}_1(K)$. The order of an element of $\text{PGL}_2(K)$ is bounded by $2 + 4[K : \mathbb{Q}]^2$, so from $2s \geq [K : \mathbb{Q}]$ it results that

$$n \leq 2 + 16s^2 < c(s). \quad (33)$$

Now we consider rational maps of degree ≥ 2 with good reduction outside S . We reduce to the hypothesis of Proposition 3. Let $(Q_{-l}, \dots, Q_{-1}, Q_0, \dots, Q_{n-1})$ be a finite orbit in $\mathbb{P}_1(K)$, for a rational map $\Psi: \mathbb{P}_1 \rightarrow \mathbb{P}_1$ defined over K with good reduction outside S , including $(Q_0, \dots, Q_{n-1})$ as a cycle for Ψ . We can associate a finite orbit in which the cycle consists of one single point (i.e. a fixed point). Indeed, the tuple $(Q_{-\lfloor \frac{l}{n} \rfloor n}, \dots, Q_{-n}, Q_0)$ is an orbit for Ψ^n and Q_0 is a fixed point. We set $m := \lfloor \frac{l}{n} \rfloor$. Of course Ψ^n can be viewed as an endomorphism of $\mathbb{P}_1$ defined over L (the extension of K defined in (6)). For every index $i \in \{0, \dots, m\}$, let $Q_{-i-n} = [t_i : l_i]$ be a representation of Q_{-i-n} in S -integral homogeneous coordinates. Recall that $\{\mathbf{a}_1, \dots, \mathbf{a}_h\}$ is a full system of representatives for the ideal classes of R_S and that the α_i 's are the S -integers verifying (5). Let $\mathbf{b}_i \in \{\mathbf{a}_1, \dots, \mathbf{a}_h\}$ be the representative of the ideal $t_i R_S + l_i R_S$. Let $\beta_i \in \{\alpha_1, \dots, \alpha_h\}$ be such that $\mathbf{b}_i^h = \beta_i R_S$. Hence there exists $\lambda_i \in K^*$ satisfying $(t_i R_S + l_i R_S)^h = \lambda_i^h \beta_i R_S$. As suggested by the referee, in L , we define

$$t'_i := \frac{t_i}{\lambda_i \sqrt[h]{\beta_i}}, \quad l'_i := \frac{l_i}{\lambda_i \sqrt[h]{\beta_i}}. \quad (34)$$

It is clear that t'_i, l'_i are elements of $\sqrt[h]{K^*}$ such that

$$(t'_i R_S + l'_i R_S) = R_S. \quad (35)$$

Furthermore, for any two distinct indices i, j

$$(t'_i l'_j - t'_j l'_i)^h = \frac{(t_i l_j - t_j l_i)^h}{\lambda_i^h \lambda_j^h \beta_i \beta_j} \in K^*.$$

By (35) with $i = 0$ there exist $r_0, s_0 \in R_S$ such that $r_0 t'_0 + s_0 l'_0 = 1$. Define the matrix

$$A = \begin{pmatrix} l'_0 & -t'_0 \\ r_0 & s_0 \end{pmatrix}$$

and further define x_i, y_i by

$$\begin{pmatrix} x_i \\ y_i \end{pmatrix} = A \begin{pmatrix} t'_i \\ l'_i \end{pmatrix}$$

for all $i \in \{0, \dots, m\}$. If we now set $P_i := [x_i : y_i]$ for all $i \in \{0, \dots, m\}$ and $\Phi := [A] \circ \Psi^n \circ [A]^{-1}$, where $[A]$ is the automorphism of $\mathbb{P}_1$ induced by A , then, by Proposition 3, $m = \left\lfloor \frac{l}{n} \right\rfloor < e^{(10)^{12}s} - 2$ and so $l < n(e^{(10)^{12}s} - 1)$. Therefore the orbit $(Q_{-l}, \dots, Q_{-1}, Q_0, \dots, Q_{n-1})$ has cardinality bounded by $ne^{(10)^{12}s}$. Since in S there are at most $s - 1$ prime ideals and $2s \geq [K : \mathbb{Q}]$, the inequality (2) becomes

$$n \leq [12(s + 1) \log(5(s + 1))]^{8s}$$

and so the theorem is proved. □

Proof of Corollary 1. Choose a finite set of generators of $\mathcal{F}$. Each of these generators has at most finitely many prime ideals of R of bad reduction. So there is a finite set S of prime ideals such that each of the chosen generators, and therefore each of the elements of $\mathcal{F}$, has good reduction outside S . We conclude by applying Theorem 1. □

References

- [1] A. T. *Introduction to Analytic Number Theory*. Springer-Verlag (1976), New York.
- [2] R. B. , *Preperiodic points of polynomials over global fields*, (2005) ArXiv:math.NT/0506480 .
- [3] F. B. H.P. S. , *The equation $x+y = 1$ in finitely generated groups*, Acta Arith. **LXXVIII.2** (1996), 189-199.
- [4] J. K. C. , *Cycles for rational maps of good reduction outside a prescribed set*, Monatsh. Math., to appear.
- [5] J. H. E. , *On equations in S -units and the Thue-Mahler equation*, Invent. Math. **75** (1984), 561-584.
- [6] J. H. E. , *The number of solutions of decomposable form equations*, Invent. Math. **122**, (1995), 559-601.
- [7] J. H. E. , H.P. S. W. M. S. , *Linear equations in variables which lie in a multiplicative group*, Ann. Math. **155**, (2002), 807-836.
- [8] S. L. , *Algebra-Revised third edition*, Springer-Verlag, GTM **211**, 2002

- [9] P. M. , J.H. S. , *Rational Periodic Points of Rational Functions*, Inter. Math. Res. Notices **2** (1994), 97-110.
- [10] P. M. , J.H. S. , *Periodic points, multiplicities, and dynamical units.*, J. Reine Angew. Math. **461** (1995), 81-122.
- [11] W. N. , *Polynomial cycles in algebraic number fields* , Colloq. Math. **58** (1989), 151-155.
- [12] W. N. , T. P. , *Finite polynomial orbits in finitely generated domain*, Monatsh. Math. **124** (1997), 309-316.
- [13] T. P. , *Polynomial cycles in certain local domains* , Acta Arith. **LXVI:1** (1994), 11-22.

Jung Kyu CANCI
 Dipartimento di Matematica e Informatica
 Università degli Studi di Udine
 via delle Scienze, 206
 33100 Udine (ITALY)
 E-mail: canci@dimi.uniud.it